

Implant Post-Op Instructions

You have just completed the first step of your implant treatment. If you look inside your mouth, you may be able to see the implant. The implant does require a clean environment in order to heal properly. In that regard, much of the success of this implant will depend on you. Proper care of your mouth will aid in the healing process and reduce the possibility of complications. The following instructions will assist you in this healing phase.

1. Following the procedure, you may find it advisable to take it easy for a day or two. Usually the fewer implants places, the sooner you will feel comfortable returning to normal activities. Jogging and vigorous activities should be avoided for a few days.
2. You may experience some swelling after the procedure. Ice packs will keep swelling and/or bruising to a minimum. Apply ice to your face on a "15 minute on a 15 minute off" cycle for the remainder of the day if possible. One way to make an effective ice pack is to put ice in a plastic bag and wrap in a towel, cloth or sock. If swelling persists after 48 hours, apply a warm moist towel.
3. You may have some discomfort after the procedure. The amount of this discomfort depends on a number of factors including the number of implants placed and where they were placed. Over the counter anti-inflammatory medicines (Advil, Aleve or Ibuprofen) are often helpful in controlling this type of discomfort. Your doctor will also prescribe a "stronger" pain medication. You may supplement the anti-inflammatory medication with this prescription medication if needed. You do not have to take or finish the pain medication.
4. Your doctor may prescribe an antibiotic for you. Unlike the pain medication, it is important that you finish this medication unless otherwise directed.
5. There is usually a minimal amount of bleeding following the procedure. We suggest that you bite on the gauze packing in your mouth for at least 20-30 minutes. If you notice bleeding from the area after the first 30 minutes, place another gauze pad over the area. Some oozing may occur for a day or two after the surgery. If implants are placed in the upper jaw, you may experience a nosebleed within the first few days after surgery.
6. Rinsing is very helpful in keeping your mouth clean. Warm salt-water rinses should be started the day after surgery. Rinse gently several times a day and continue to do so for ten to fourteen days.
7. Do not attempt to floss, brush or water-pik in the area of surgery for the first two weeks. Keep your mouth clean with warm water rinses. Following the healing period, you will be instructed on further cleaning techniques. The rest of your mouth should continue to be cleaned as usual.
8. Diet is very important in the postoperative period. We would like you to eat soft nutritious foods. Care should be taken to avoid hard or crunchy foods that might cause trauma to the implant area. Gently rinse with warm water after every meal.
9. Smoking should be avoided for at least two weeks following surgery. This has been shown to negatively affect the healing process.
10. Your doctor will decide when dentures or partial dentures may be worn after surgery, but they must be worn only with caution. Remember, no undue pressure should be applied over that implant surgical site. This can affect the success of your implant.
11. Stitches will be removed in 7-10 days, as deemed necessary by the doctor.
12. Should you notice any swelling, redness or discomfort in the area of the implant after the healing phase, PLEASE NOTIFY THE OFFICE. Your patience and self-care will help make your dental implant successful. If you have any questions or unusual problems, please call us at any time 604-267-3368 or Dr. Benny Yee cell: 604-619-1989.